


NAVIGATING SCALING: MODELLING AND ANALYSING

P. Abry⁽¹⁾,

P. Gonçalves⁽²⁾

(1) SISYPH, CNRS,
Ecole Normale Supérieure
Lyon, France

(2) INRIA Rhône-Alpes,
On Leave at IST-ISR, Lisbon




IN COLLABORATIONS WITH :

P. Flandrin, D. Veitch,
P. Chainais, B. Lashermes, N. Hohn,
S. Roux, P. Borgnat,
M. Taqqu, V. Pipiras, R. Riedi, S. Jaffard

Wavelet And Multifractal Analysis, Cargèse, France, July 2004.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 07 JAN 2005		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Navigating Scaling: Modelling And Analysing				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) SISYPH, CNRS, Ecole Normale Sup'erieure Lyon, France				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001750, Wavelets and Multifractal Analysis (WAMA) Workshop held on 19-31 July 2004., The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 64	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

SCALING PHENOMENA ?



- **DETECTION:** SCALING ? WHAT DOES IT MEAN ? NON STATIONARITY ?
- **IDENTIFICATION:** RELEVANT STOCHASTIC MODELS ?
- **ESTIMATION:** RELEVANT PARAMETER ESTIMATION ?
- **SIDE ISSUES:**
ROBUSTNESS ? COMPUTATIONAL COST ? REAL TIME ? ON LINE ?

OUTLINE

I. INTUITIONS, MODELS, TOOLS

- I.1 INTUITIONS, DEFINITION, APPLICATIONS
- I.2 STOCHASTIC MODELS: SELF-SIMILARITY VS MULTIFRACTAL
- I.3 MULTIREOLUTION TOOLS, AGGREGATION, INCREMENTS
- I.4 WAVELETS, CONTINUOUS, DISCRETE

II. SECOND ORDER ANALYSIS, SELF SIMILARITY AND LONG MEMORY





- II.1 RANDOM WALKS, SELF SIMILARITY, LONG MEMORY,
- II.2 2ND ORDER WAVELET STATISTICAL ANALYSIS,
- II.3 ESTIMATION, ESTIMATION PERFORMANCE,
- II.4 ROBUSTNESS AGAINST NON STATIONARITIES,

III. HIGHER ORDER ANALYSIS, MULTIFRACTAL PROCESSES

- III.1 MULTIPLICATIVE CASCADES, MULTIFRACTAL PROCESSES,
- III.2 HIGHER ORDER WAVELET STATISTICAL ANALYSIS,
- III.3 FINITENESS OF MOMENTS,
- III.4 ESTIMATION, ESTIMATION PERFORMANCE,
- III.5 NEGATIVE ORDERS,
- III.6 BEYOND POWER LAWS.


IRREGULARITIES, VARIABILITIES

SCALING OR NON STATIONARITIES?




SCALING ?


Traffic (WAN) Internet




Traffic (WAN) Internet



SCALING ?



SCALING ?





SCALING !

- **DEFINITION :**







NON PROPERTY: NO CHARACTERISTIC SCALE.
NON GAUSSIAN, NON STATIONARY, NON LINEAR

- **EVIDENCE:**

The whole resembles to its part, the part resembles to the whole.



- **ANALYSIS:** Rather than for a characteristic scale, look for a relation, a mechanism, a cascade between scales.



SCALING : OPERATIONAL DEFINITIONS

- MULTIRESOLUTION QUANTITY:

$$T_X(a, t) \quad (\text{e.g., Wavelet Coef.}).$$

- POWER LAWS:

$$\begin{aligned} \mathbf{E}|T_X(a, t)|^q &= c_q |a|^{\zeta(q)}, \\ \frac{1}{n} \sum_{k=1}^n |T_X(a, t_k)|^q &= c_q |a|^{\zeta(q)}, \end{aligned}$$





- FOR A RANGE OF SCALES a ,
- FOR A RANGE OF ORDERS q
- SCALING EXPONENTS $\zeta(q)$.

- BEYOND POWER LAWS : WARPED INF. DIV. CASCADES

$$\begin{aligned} \mathbf{E}|T_X(a, t)|^q &= C_q |a|^{\zeta(q)} = C_q \exp(\zeta(q) \ln a) \\ \mathbf{E}|T_X(a, t)|^q &= C_q \exp(\zeta(q) n(a)) \end{aligned}$$

→ VISIT PIERRE CHAINAIS'S POSTER

UBIQUITY ?



UBIQUITY !


- Hydrodynamic Turbulence,
- Physiology, Biological Rhythms (Heart beat, walk),
- Geophysics (Faults Repartition, Earthquakes),
- Hydrology (Water Levels),
- Statistical Physics (Long Range Interactions),
- Thermal Noises (semi-conductors),
- Information Flux on Networks, Computer Network Traffic,
- Population Repartition (local: cities, global: continent),
- Financial Markets (Daily returns, Volatility, Currencies Exchange Rates),
- ...

ANALYSING TOOL 1 : AGGREGATION

COMPARE DATA AGAINST A BOX, THEN VARY a

$$T_X(a, t) = \frac{1}{aT_0} \int_t^{t+aT_0} X(u) du$$

AVERAGE




WORKS ONLY FOR POSITIVE TIME SERIES, DENSITY

ANALYSING TOOL 2 : INCREMENTS

COMPARE DATA AGAINST A DIFFERENCE OF DELTA FUNCTIONS, THEN VARY a

$$T_X(a, t) = X(t + a\tau_0) - X(t)$$

DIFFERENCE



INCREMENTS OF HIGHER ORDERS OR GENERALISED N -VARIATIONS

- Order 2 : $T_X(a, t) = -X(t + 2a\tau_0) + 2X(t + a\tau_0) - X(t)$,
 - Order N : $T_X(a, t) = \sum_{p=0}^N (-1)^p a_p X(t + pa\tau_0)$,
- where $\sum_{p=0}^N (-1)^p a_p p^k \equiv 0, \quad k = 0, \dots, N - 1$.

ANALYSING TOOL: MULTIREOLUTION ANALYSIS

• MULTIREOLUTION QUANTITIES:


$$X(t) \rightarrow T_X(a, t) = \langle f_{a,t} | X \rangle, \quad f_{a,t}(u) = \frac{1}{a} f_0\left(\frac{u-t}{a}\right)$$

AGGREGATION

$$f_0(u) = (\beta_0)$$

$$= \frac{1}{aT_0} \int_t^{t+aT_0} X(u) du$$

BOX, AVERAGE




INCREMENTS

$$f_0(u) = (I_0)$$

$$= X(t + a\tau_0) - X(t)$$

DIFFERENCE



?

?

?

?

?

ANALYSING TOOL: MULTIREOLUTION ANALYSIS

- MULTIREOLUTION QUANTITIES:

$$X(t) \rightarrow T_X(a, t) = \langle f_{a,t} | X \rangle, \quad f_{a,t}(u) = \frac{1}{a} f_0\left(\frac{u-t}{a}\right)$$


- CHOICES FOR MOTHER FUNCTIONS: f_0 ,

AGGREGATION

$$f_0(u) = (\beta_0)^{*N}$$

$$= \frac{1}{aT_0} \int_t^{t+aT_0} X(u) du$$

BOX, AVERAGE




N

INCREMENTS

$$f_0(u) = (I_0)^{*N}$$

$$= X(t + a\tau_0) - X(t)$$

DIFFERENCE




N

WAVELETS

$$f_0(u) = \psi_{0,N}$$

$$= \int X(u) \frac{1}{|a|} \psi_0\left(\frac{u-t}{a}\right),$$




AVERAGE, DIFFERENCE






N

WAVELETS AND SCALING: KEY INGREDIENTS

- DILATION OPERATOR, $\frac{1}{|a|}\psi_0\left(\frac{t}{|a|}\right)$




- NUMBER OF VANISHING MOMENTS,
$$N \geq 1, \int t^k \psi_0(t) dt \equiv 0, \quad k = 0, 1, \dots, N-1.$$



WAVELET TRANSFORMS

- MOTHER-WAVELET AND "BASIS": $\int \psi_0(u)du = 0, \quad \psi_{a,t}(u) = \frac{1}{|a|}\psi_0\left(\frac{u-t}{a}\right)$
- WAVELET COEFFICIENTS: CONTINUOUS WT $T_X(a, t) = \langle X, \psi_{a,t} \rangle$


SCALE



TIME

MODULUS MAXIMA WT


SKELETON : MAXIMA LINES



AND DISCRETE WT

$$d_X(j, k) = T_X(a = 2^j, t = 2^j k)$$

SCALE



TIME

OUTLINE

I. INTUITIONS, MODELS, TOOLS

- I.1 INTUITIONS, DEFINITION, APPLICATIONS
- I.2 STOCHASTIC MODELS: SELF-SIMILARITY VS MULTIFRACTAL
- I.3 MULTIREOLUTION TOOLS, AGGREGATION, INCREMENTS
- I.4 WAVELETS, CONTINUOUS, DISCRETE

II. SECOND ORDER ANALYSIS, SELF SIMILARITY AND LONG MEMORY

- II.1 RANDOM WAKS, SELF SIMILARITY, LONG MEMORY,
- II.2 2ND ORDER WAVELET STATISTICAL ANALYSIS,
- II.3 ESTIMATION, ESTIMATION PERFORMANCE,
- II.4 ROBUSTNESS AGAINST NON STATIONARITIES,

III. HIGHER ORDER ANALYSIS, MULTIFRACTAL PROCESSES

- III.1 MULTIPLICATIVE CASCADES, MULTIFRACTAL PROCESSES,
- III.2 HIGHER ORDER WAVELET STATISTICAL ANALYSIS,
- III.3 FINITENESS OF MOMENTS,
- III.4 ESTIMATION, ESTIMATION PERFORMANCE,
- III.5 NEGATIVE ORDERS,
- III.6 BEYOND POWER LAWS.

MOD. TOOL 1: RAND. WALKS AND SELF SIMILARITY

RANDOM WALK: $X(t + \tau) = X(t) + \underbrace{\delta_\tau X(t)}_{\text{Steps or Increments}}$

STATISTICAL PROPERTIES OF THE STEPS:

- **A1:** Stationary,
- **A2:** Independent,
- **A3:** Gaussian,
 - \Rightarrow Ordinary Random Walk, Ordinary Brownian Motion,
 - $\Rightarrow \mathbf{E}X(t)^2 = 2D|t|$, Einstein relation,
 - $\Rightarrow \mathbf{E}|X(t)|^q = 2D|t|^{q/2}$, $q > -1$.

ANOMALIES:

- $\Rightarrow \mathbf{E}X(t)^2 = 2D|t|^\gamma$,
- $\Rightarrow \mathbf{E}X(t)^2 = \infty$.

SELF SIMILAR RANDOM WALKS:


- **B1:** Stationary,
- **B2:** Self Similarity

MODELLING TOOL 1: SELF-SIMILARITY

- **DEFINITION:** $\delta_\tau X(t) \stackrel{fdd}{=} c^H \delta_{\tau/c} X(t/c), \forall c > 0$, DILATION FACTOR,
 $1 > H > 0$: SELF-SIMILARITY EXPONENT

- **INTERPRETATIONS:**

- COVARIANCE UNDER DILATION (CHANGE OF SCALE),
- THE WHOLE AND THE SUBPART (STATISTICALLY) UNDISTINGUISHABLE,
- NO CHARACTERISTIC SCALE OF TIME.



- **IMPLICATIONS:**

- NON STATIONARITY PROCESS WITH STATIONARY INCREMENTS
- $\mathbb{E}|X(t + a\tau_0) - X(t)|^q = C_q |a|^{qH}$,
- $\forall a > 0, \forall c > 0, \forall q \quad \mathbb{E}|X(t)|^q < \infty$,
- A SINGLE SCALING EXPONENT H .
- ADDITIVE STRUCTURE,
- (CORRELATED) RANDOM WALK, LONG MEMORY, LONG RANGE CORRELATIONS.

MOD. TOOL 1 (BIS): LONG RANGE DEPENDENCE

● DEFINITIONS :

- LET X BE A 2ND STATIONARY PROCESS WITH,
- COVARIANCE : $c_X(\tau) = \mathbb{E}X(t)X(t + \tau)$
- SPECTRUM : $\Gamma_X(\nu)$

$$\begin{aligned}c_X(\tau) &= c_\tau |\tau|^{-\beta}, & 0 < \beta < 1, & \quad |\tau| \rightarrow +\infty \\ \Gamma_X(\nu) &= c_f |\nu|^{-\alpha}, & 0 < \alpha < 1, & \quad |\nu| \rightarrow 0\end{aligned}$$

WITH $\alpha = 1 - \beta$ AND $c_f = 2(2\pi) \sin((1 - \gamma)\pi/2)c_\tau$.

● CONSEQUENCES :

- $\sum_A^{+\infty} c_X(\tau) d\tau = +\infty, A > 0,$
- NO CHARACTERISTIC SCALE,
- AGGREGATION : $T_X(\mathbf{a}, t) = \frac{1}{\mathbf{a}T_0} \int_t^{t+\mathbf{a}T_0} X(u) du,$
 $\Rightarrow \text{VAR } T_X(\mathbf{a}, t) \sim C \mathbf{a}^{\alpha-1}, \mathbf{a} \rightarrow +\infty,$
- INCREMENTS OF SELF.-SIM. PROC. (WITH $H > 1/2$)
ARE LONG RANGE DEP. (WITH $\alpha = 2H - 1$).

WAVELETS AND SELF-SIMILAR PROCESSES WITH STATIONARY INCREMENTS - SUMMARY

(Flandrin et al., Tewfik and Kim)

• **P1:** $\{d_X(j, k), k \in \mathbb{Z}\}$ **STATIONARY** Sequences for each Scale 2^j .
 $N \geq 1$

• **P2:** **SELF-SIMILARITY** : **Dilation**


$$\{X(t)\} \stackrel{d}{=} \{c^H X(t/c)\} \Rightarrow \{d_X(0, k)\} \stackrel{d}{=} \{2^{-jH} d_X(j, k)\}$$

- **P3 :** **MARGINAL DIST.** $P_j(d) = \frac{1}{\beta_0} P_{j'}(\frac{d}{\beta_0}), \quad \beta_0 = \left(\frac{2^{j'}}{2^j}\right)^H$.

• **P4 :** $\{d_X(j, k)\}$ **SHORT RANGE DEPENDENT** IF $N > H + 1/2$.
 $|2^j k - 2^{j'} k'| \rightarrow +\infty, \quad |\text{Cov } d_X(j, k) d_X(j', l)| \leq D |2^j k - 2^{j'} k'|^{2(H-N)},$
 $N \geq 1$ and **Dilation**

WAVELETS AND LONG RANGE DEPENDENCE

(Flandrin)



WAVELETS AND SELF-SIMILAR PROCESSES WITH STATIONARY INCREMENTS - SUMMARY

• **P1:** $\{d_X(j, k), k \in \mathbb{Z}\}$ **STATIONARY** Sequences for each Scale 2^j .
 $N \geq 1$

• **P2:** **SELF-SIMILARITY : Dilation**

$$\{X(t)\} \stackrel{d}{=} \{c^H X(t/c)\} \Rightarrow \{d_X(0, k)\} \stackrel{d}{=} \{2^{-jH} d_X(j, k)\}$$

- **P3 :** **MARGINAL DIST.** $P_j(d) = \frac{1}{\beta_0} P_{j'}(\frac{d}{\beta_0})$, $\beta_0 = \left(\frac{2^{j'}}{2^j}\right)^H$.

• **P4 :** $\{d_X(j, k)\}$ **SHORT RANGE DEPENDENT** IF $N > H + 1/2$.
 $|2^j k - 2^{j'} k'| \rightarrow +\infty$, $|\text{Cov } d_X(j, k) d_X(j', k')| \leq D |2^j k - 2^{j'} k'|^{2(H-N)}$,
 $N \geq 1$ and **Dilation**

\Rightarrow **IDEALISATION :** $d_X(j, k)$ **INDEPENDENT VARIABLES**.

\Rightarrow **INTERPRETATIONS:** $X(t) = \sum_k a_X(J, k) \varphi_{J,k}(t) + \sum_{j=1, \dots, J, k} d_X(j, k) \psi_{j,k}(t)$.

\Rightarrow **IMPLICATIONS:** $\mathbf{E}|d_X(j, k)|^q = \mathbf{E}|d_X(0, k)|^q 2^{\mathbf{j} q H}$ $\forall q / \mathbf{E}|d_X(0, k)|^q < \infty$.

WAVELETS AND LONG RANGE DEPENDENCE

- **SPECTRAL ANALYSIS :**

Let X be a 2nd Order stationary process,

Let Ψ be the FT of ψ with central frequency ν_0 and bandwidth $\Delta\nu_0$.

$$\begin{aligned}\mathbf{E}|d_X(j, k)|^2 &= \int \Gamma_X(\nu) |\Psi(2^j \nu)| d\nu \\ &\simeq 2^{-j} \Gamma_X(2^{-j} \nu_0) \text{ within bandwidth } 2^{-j} \Delta\nu_0.\end{aligned}$$

- **LET X BE LONG RANGE DEPENDENT :**

- **POWER LAW:** $\Gamma_X(\nu) = c_f |\nu|^{-\alpha}, 0 < \alpha < 1, |\nu| \rightarrow 0$

- **POWER LAW:** $\mathbf{E}|d_X(j, k)|^2 \sim C 2^{j(\alpha-1)}, j \rightarrow +\infty,$

- $\{d_X(j, k)\}$ **SHORT RANGE DEPENDENT** IF $N > \alpha - 1$.

$|2^j k - 2^{j'} k'| \rightarrow +\infty, |\text{Cov } d_X(j, k) d_X(j', k')| \leq D |2^j k - 2^{j'} k'|^{\alpha-1-2N},$

$N \geq 1$ and Dilation

2ND ORDER WAVELET STATISTICAL ANALYSIS


Abry, Gonçalves, Flandrin

PRINCIPLES:


- IDEAS : **P1** $\Rightarrow \mathbf{E}|d_X(j, k)|^2 = C_2 2^{j2H}$
 $\Rightarrow \log_2 \mathbf{E}|d_X(j, k)|^2 = j2H + \beta_q,$
- PROBLEMS: ESTIMATE $\mathbf{E}|d_X(j, k)|^2$ FROM A SINGLE FINITE LENGTH OBSERVATION ?
- SOLUTION : **P2 et P3** \Rightarrow STATISTICAL AVERAGES \Rightarrow TIME AVERAGES,
 $S_2(j) = (1/n_j) \sum_{k=1}^{n_j} |d_X(j, k)|^2$

LOG-SCALE DIAGRAMS: $\log_2 S_2(j)$ vs $\log_2 2^j = j$

2ND ORDER WAVELET-BASED STATISTICAL ANALYSIS FOR SELF-SIMILARITY




2ND ORDER WAVELET-BASED STATISTICAL ANALYSIS FOR LONG RANGE DEPENDENCE



WAVELETS AND 2ND-ORDER SCALING: ESTIMATION

- **DYADIC GRID** (DISCRETE WAVELET TRANSFORM): $a_j = 2^j, t_{j,k} = k2^j,$



- **STRUCTURE FUNCTION (TIME AVERAGE):**

$$Y_j = (\frac{1}{2} \log_2 S_2(2^j)) = \frac{1}{2} \log_2(1/n_j) \sum_{k=1}^{n_j} |d_X(j, k)|^2$$

- **DEFINITION :**

$$Y_j \text{ versus } \log_2 2^j = j, \\ \hat{H} = \sum_{j=j_1}^{j_2} w_j Y_j .$$

$$\text{WHERE } \sum_j j w_j \equiv 1, \quad \sum_j w_j \equiv 0, \text{ WITH } w_j \equiv \frac{B_0 j - B_1}{B_0 B_2 - B_1^2}, \\ \text{AND } p = 0, 1, 2, \quad B_p = \sum_j j^p / a_j, \quad a_j \text{ ARBITRARY NUMBERS.}$$

- **WHAT ARE THE PERFORMANCE OF SUCH AN ESTIMATOR ?**
WHEN APPLIED TO A SELF-SIMILAR. OR LRD PROCESS

WAVELETS AND 2ND-ORDER SCALING: ESTIMATION

Abry, Gonçalves, Flandrin,

Abry, Veitch


- **ASSUME:** - *i*) X GAUSSIAN,
- *ii*) IDEALISATION: EXACT INDEPENDENCE.
- **BIAS :**
$$\mathbf{E} \log_2 S_2(j) = \log_2 \mathbf{E} S_2(j) + \underbrace{\Gamma'(n_j/2) - \log_2(n_j/2)}_{g_j}.$$
$$\Rightarrow \mathbf{E} \hat{H} = H + \frac{1}{2} \sum_j w_j g_j.$$
- **VARIANCE:**
 - $\text{Var } \hat{H} = \frac{1}{4} \sum_j w_j^2 \sigma_j^2,$
 - $\min \text{Var } \hat{H} \Rightarrow a_j \propto \text{Var } \log_2 S_2(j)$
 - $\text{Var } \log_2 S_2(j) \simeq C/n_j \simeq 2^j C/n,$
$$\Rightarrow \text{VAR } \hat{H} \simeq \left((\log_2 e)^2 (\sum_j w_j^2 2^j) \right) / \textcolor{red}{n},$$
$$\Rightarrow \text{ANALYTICAL (APPROXIMATE) CONFIDENCE INTERVAL}$$
$$(\text{DOES NOT DEPEND ON UNKNOWN } H).$$
- **ACTUAL PERFORMANCES :** NEGLIGIBLE BIAIS, EXTREMELY CLOSE TO MLE.
- **CONCEPTUAL AND PRACTICAL SIMPLICITY :** MATLAB CODE AVAILABLE.

WAV. AND 2ND-ORDER SCALING: ROBUSTNESS

Superimposed Trends

$$Y(t) = X(t) + T(t) \Rightarrow d_Y(j, k) = d_X(j, k) + d_T(j, k)$$


- If $T(t)$ Polynomial of degree P , then $d_T \equiv 0$ when $N > P$,
- If $T(t)$ smooth trend, then the d_T decrease as N increases.



Vary N !





WAV. AND 2ND-ORDER SCALING: ROBUSTNESS

Superimposed Trends - Ethernet Data (Veitch, Abry)



WAV. AND 2ND-ORDER SCALING: ROBUSTNESS

Constancy along time of Scaling laws (Veitch, Abry)



SELF-SIMILARITY

- SELF-SIMILARITY:

$$\mathbf{E}|d_X(j, k)|^q = C_q(2^j)^{qH}$$

- Power Laws,
- $\forall 2^j$ (for all scales),
- $\forall q / \mathbf{E}|d_X(j, k)|^q < \infty$,
- A single parameter H
- Additive Structure.

- ?

- ?

BEYOND SELF-SIMILARITY

- SELF-SIMILARITY:

$$\mathbf{E}|d_X(j, k)|^q = C_q(2^j)^{qH}$$

- Power Laws,
- $\forall 2^j$ (for all scales),
- $\forall q / \mathbf{E}|d_X(j, k)|^q < \infty$,
- A single parameter H
- Additive Structure.

- MULTIFRACTAL

$$\mathbf{E}|d_X(j, k)|^q = C_q(2^j)^{\zeta(q)}$$

- Power Laws,
- $\forall 2^j < L$, (for fine scales only, in the limit $2^j \rightarrow 0$),
- $\forall q$?
- A whole collection of scaling parameter $\zeta(q)$
- Multiplicative Structure.

- ?

OUTLINE

I. INTUITIONS, MODELS, TOOLS

- I.1 INTUITIONS, DEFINITION, APPLICATIONS
- I.2 STOCHASTIC MODELS: SELF-SIMILARITY VS MULTIFRACTAL
- I.3 MULTIREOLUTION TOOLS, AGGREGATION, INCREMENTS
- I.4 WAVELETS, CONTINUOUS, DISCRETE

II. SECOND ORDER ANALYSIS, SELF SIMILARITY AND LONG MEMORY

- II.1 RANDOM WALKS, SELF SIMILARITY, LONG MEMORY,
- II.2 2ND ORDER WAVELET STATISTICAL ANALYSIS,
- II.3 ESTIMATION, ESTIMATION PERFORMANCE,
- II.4 ROBUSTNESS AGAINST NON STATIONARITIES,


III. HIGHER ORDER ANALYSIS, MULTIFRACTAL PROCESSES

- III.1 MULTIPLICATIVE CASCADES, MULTIFRACTAL PROCESSES,
- III.2 HIGHER ORDER WAVELET STATISTICAL ANALYSIS,
- III.3 FINITENESS OF MOMENTS,
- III.4 ESTIMATION, ESTIMATION PERFORMANCE,
- III.5 NEGATIVE ORDERS,
- III.6 BEYOND POWER LAWS.

MODELLING TOOL 2: MULTIPLICATIVE CASCADES

● DEFINITION:

- SPLIT DYADIC INTERVALS $I_{j,k}$ INTO TWO,
- I.I.D. MULTIPLIERS $W_{j,k}$
- $Q_J(t) = \prod_{\{(j,k): 1 \leq j \leq J, t \in I_{j,k}\}} W_{j,k}$,



● IMPLICATIONS:


- LOCAL HOLDER EXPONENT,
- MULTIFRACTAL SAMPLE PATHS, MULTIFRACTAL SPECTRUM $D(h)$
- CASCADES, MULTIPLICATIVE STRUCTURE,
- $\sum_k \left(1/a \int_{t_k}^{t_k + a\tau_0} X(u) du \right)^q = C_q |a|^{\zeta_q}$, FINE SCALES $a \rightarrow 0$,
- MULTIPLE EXPONENTS ζ_q ,
- NO CHARACTERISTIC SCALE,
- $\zeta_q = -\log_2 \mathbb{E} W^q$, NON LINEAR IN q .

MODELLING TOOL 2: MULTIPLICATIVE CASCADES

YAGLOM, MANDELBROT

MANDELBROT'S CASCADE (CMC)


- IID W ,
- DYADIC GRID,



BARRAL, MANDELBROT

COMPOUND POISSON CASCADE (CPC)


- IID W ,
- POINT PROCESS,



SCHMITT ET AL.,
BACRY ET AL., CHAINAIS ET AL.

INFINITELY DIVISIBLE CASCADE (IDC)

- CONTINUOUS INFINITELY
- DIVISIBLE MEASURE M ,



$$Q_r(t) = \prod W_{j,k}, \quad \prod W_{j,k}, \quad \exp \int dM(t', r'),$$

$$\varphi(q) = -\log_2 \mathbb{E} W^q, \quad = -q(1 - \mathbb{E} W) + 1 - \mathbb{E} W^q, \quad = \rho(q) - q\rho(1),$$


$$A(t) = \lim_{r \rightarrow 0} \int_0^t Q_r(u) du,$$

FOR A RANGE OF q S, $\mathbb{E} |A(t + \textcolor{red}{a}\tau_0) - A(t)|^{\textcolor{blue}{q}} = c_q |\textcolor{red}{a}|^{\textcolor{blue}{q} + \varphi(q)}$,
RESOLUTION DEPTH < SCALE < INTEGRAL SCALE, $a_m = r < \textcolor{red}{a} < a_M = L$.

MULTIFRACTAL PROCESSES


DENSITY: $Q_r(t) = \Pi W_{j,k}$

$$\mathbb{E} \left(\frac{1}{a} \int_t^{t+a\tau_0} Q_r(u) du \right)^q = c_q a^{\varphi(q)},$$



MEASURE: $A(t) = \lim_{r \rightarrow 0} \int_0^t Q_r(u) du,$


$$\mathbb{E} |A(t + a\tau_0) - A(t)|^q = c_q |a|^{qH + \varphi(qH)},$$



**FRACTIONAL BROWNIAN MOTION
IN MULTIFRACTAL TIME:**

$$V_H(t) = B_H(A(t)),$$


$$\mathbb{E} |V_H(t + a\tau_0) - V_H(t)|^q = c_q |a|^{qH + \varphi(qH)},$$



MULTIFRACTAL RANDOM WALK:

$$Y_H(t) = \int_0^t Q_r(s) dB_H(s),$$

$$\mathbb{E} |Y_H(t + a\tau_0) - Y_H(t)|^q = c_q |a|^{qH + \varphi(q)}.$$



MATLAB SYNTHESIS ROUTINES : CHAINAIS, ABRY




HIGHER-ORDER WAVELET STATISTICAL ANALYSIS

PRINCIPLES :

- IDEAS : **P1** $\Rightarrow \mathbb{E}|d_X(j, k)|^q = \mathbb{E}|d_X(0, k)|^q 2^{j\zeta_q}$
 $\Rightarrow \log_2 \mathbb{E}|d_X(j, k)|^q = j\zeta_q + \beta_q,$
- PROBLEMS: ESTIMATE $\mathbb{E}|d_X(j, k)|^q$ FROM A SINGLE FINITE LENGTH OBSERVATION ?
- SOLUTION : **P2 et P3** \Rightarrow STATISTICAL AVERAGES \Rightarrow TIME AVERAGES,
 $S_q(j) = (1/n_j) \sum_{k=1}^{n_j} |d_X(j, k)|^q$


LOG-SCALE DIAGRAMS: $\log_2 S_q(j)$ vs $\log_2 2^j = j$

LOGSCALE DIAGRAMS - MULTIFRACTAL PROC.



WAV. AND HIGHER-ORDER SCALING: ESTIMATION

- DYADIC GRID (DISCRETE WAVELET TRANSFORM): $a_j = 2^j, t_{j,k} = k2^j,$



- STRUCTURE FUNCTIONS (TIME AVERAGE):

$$S_q(j) = (1/n_j) \sum_{k=1}^{n_j} |d_X(j, k)|^q$$

- DEFINITION:

$$Y_{j,q,n} = \log_2 S_n(2^j, q; f_0) \text{ VERSUS } \log_2 2^j = j,$$

$$\hat{\zeta}(q, n) = \sum_{j=j_1}^{j_2} w_{j,q} Y_{j,q,n} .$$

NON WEIGTHED: $a_j = cste$

- WHAT ARE THE PERFORMANCE OF SUCH ESTIMATORS ?
WHEN APPLIED TO MULTIFRACTAL PROCESSES

TEST FOR THE FINITENESS OF MOMENTS

GONÇALVÈS, RIEDI

THEOREM :

LET \mathbf{X} BE A RV WITH CHARACTERISTIC FUNCTION $\chi(s) := \mathbb{E} \exp\{is\mathbf{X}\}$.

IF $\mathcal{H}_{\Re\chi} := \sup\{\alpha > 0 : |\Re\chi(s) - P_\alpha(s)| \leq C|s|^\alpha\}$,

IS THE LOCAL HÖLDER REGULARITY OF $\Re\chi$ AT THE ORIGIN, THEN

$\mathbb{E}|\mathbf{X}|^q < +\infty \forall q \leq q_c^+$ AND $\mathcal{H}_{\Re\chi} \leq q_c^+ \leq \lfloor \mathcal{H}_{\Re\chi} \rfloor + 1$.


ESTIMATOR :

$\{X_k\}_{k=1,\dots,n}$, n I.I.D RVs, SET





$$W(a) := n^{-1} \sum_{k=1}^n \Psi(a \cdot X_k)$$

WITH Ψ A REAL AND SEMI-DEFINITE
FOURIER TRANSFORM OF A
SUFFICIENTLY REGULAR WAVELET ψ .
THEN

$$\mathcal{H}_{\Re\chi} = \limsup_{a \rightarrow 0^+} \frac{\log |W(a)|}{\log a}.$$




ESTIMATING THE PARTITION FUNCTION SUPPORT



METHODOLOGY

- **NUMERICAL SYNTHESIS OF PROCESSES:**
 - ACCUMULATE n_{real} NUMERICAL REPLICATIONS WITH LENGTH n SAMPLES.
- **APPLY SCALING EXPONENTS ESTIMATORS:**
 - COMPUTE $\hat{\zeta}(q, n)_{(l)}$ FOR EACH REPLICATION,
 - AVERAGE OVER REPL. TO OBTAIN THE STATISTICAL PERFORMANCE OF $\hat{\zeta}(q, n)$
- **ASYMPTOTIC BEHAVIOURS:**
 - THE CASCADE DEPTH INCREASES FOR A GIVEN NUMBER OF INTEGRAL SCALES.
 - ... ,



METHODOLOGY

- **NUMERICAL SYNTHESIS OF PROCESSES:**


- ACCUMULATE n_{real} NUMERICAL REPLICATIONS WITH LENGTH n SAMPLES.

- **APPLY SCALING EXPONENTS ESTIMATORS:**

- COMPUTE $\hat{\zeta}(q, n)_{(l)}$ FOR EACH REPLICATION,
- AVERAGE OVER REPL. TO OBTAIN THE STATISTICAL PERFORMANCE OF $\hat{\zeta}(q, n)$

- **ASYMPTOTIC BEHAVIOURS:**


- THE CASCADE DEPTH INCREASES FOR A GIVEN NUMBER OF INTEGRAL SCALES.
- THE NUMBER OF INTEGRAL SCALES INCREASES FOR A GIVEN CASCADE DEPTH,




LINEARISATION EFFECT: $\hat{\zeta}(q)$

LASHERMES, ABRY, CHAINAIS

CPC $Q_r EI(1)$



CPC $V_H EIII(3)$




$q > q_o, \hat{\zeta}(q, n) = \alpha_o + \beta_o q, \quad q_o, \alpha_o, \beta_o \text{ ARE RV.}$


LINEARISATION EFFECT: LEGENDRE TRANSFORM

$$D(h) = d + \min_q(qh - \zeta(q)), \text{ (} d \text{ EUCLIDIEN DIMENSION OF SPACE).}$$

CPC Q_r $EI(1)$




CPC V_H $EIII(3)$




ACCUMULATION POINTS : $D_o(h_o)$, WITH $D_o = d - \alpha_o$, $h_o = \beta_o$,
 D_o, h_o ARE RV.

LIN. EFFECT: ASYMPTOTIC BEHAVIOURS

- GIVEN RESOLUTION, INCREASING NUMBER OF INTEGRAL SCALES,



- GIVEN NUMBER OF INTEGRAL SCALES, INCREASING RESOLUTION,



LINEARISATION EFFECT: CONJECTURE

- CRITICAL POINTS:


$$\begin{cases} D_*^\pm &= 0, \\ D(h_*^\pm) &= 0, \\ h_*^\pm &= (d\zeta(q)/dq)_{q=q_*^\pm}. \end{cases}$$

- RESULTS:

$$EI : \begin{cases} \hat{\zeta}(q, n) = d - D_o^- + h_o^- q & \rightarrow d - D_*^- + h_*^- q, & q \leq q_*^-, \\ \hat{\zeta}(q, n) & \rightarrow \zeta(q), & q_*^- \leq q \leq q_*^+, \\ \hat{\zeta}(q, n) = d - D_o^+ + h_o^+ q & \rightarrow d - D_*^+ + h_*^+ q, & q_*^+ \leq q. \end{cases}$$

$$EII \& III : \begin{cases} \hat{\zeta}(q, n) & \rightarrow \zeta(q), & 0 < q \leq q_*^+, \\ \hat{\zeta}(q, n) = d - D_o^+ + h_o^+ q & \rightarrow d - D_*^+ + h_*^+ q, & q_*^+ \leq q. \end{cases}$$

- ILLUSTRATION:




LINEARISATION EFFECT: COMMENTS


WHEN DOES THE LINEARISATION EFFECT EXIST ?

- FOR ALL TYPES OF CASCADES: CMC, CPC, IDC,
- FOR ALL TYPES OF PROCESSES: Q_r, A, V_H, Y_H ,
- FOR ALL NUMBERS OF VANISHING MOMENTS: $N \geq 1$,
- FOR ALL MRA-BASED ESTIMATORS: WAVELETS, INCREMENTS, AGGREGATION,
- CAN BE WORKED OUT FOR $q < 0$,
- EXTENDS TO DIMENSION HIGHER THAN $d > 1$.

EXTENSION: STANDARD WT VERSUS WTMM (1/3).





EXTENSION: 2D MULTIPLICATIVE CASCADE (2/3).



EXTENSION: 3D MULTIPLICATIVE CASCADE (3/3).

3D CMC (LOG NORMAL), EI(1) COMPARED TO A 2D SLICE.



LINEARISATION EFFECT: COMMENTS

WHEN DOES THE LINEARISATION EFFECT EXIST ?

- FOR ALL TYPES OF CASCADES: CMC, CPC, IDC,
- FOR ALL TYPES OF PROCESSES: Q_r, A, V_H, Y_H ,
- FOR ALL NUMBERS OF VANISHING MOMENTS: $N \geq 1$,
- FOR ALL MRA-BASED ESTIMATORS: WAVELETS, INCREMENTS, AGGREGATION,
- CAN BE WORKED OUT FOR $q < 0$,
- EXTENDS TO DIMENSION HIGHER THAN $d > 1$.

WHAT THE LINEARISATION EFFECT IS NOT:

- A LOW PERFORMANCE ESTIMATION EFFECT.
- A FINITE SIZE EFFECT : THE CRITICAL PARAMETERS DO NOT DEPEND ON n ,
BE IT THE NUMBER OF INTEGRAL SCALES,
OR THE DEPTH (OR RESOLUTION) OF THE CASCADES.
- A FINITENESS OF MOMENTS EFFECT,
 - $q_c^- < 0 < 1 < q_c^+$, $q - 1 + \varphi(q) = 0$,
 - $q_c^- < q_*^- < 0 < 1 < q_*^+ < q_c^+$,

WHAT THE LINEARISATION EFFECT MIGHT BE:

- MULTIPLICATIVE MARTINGALES ?
- OSSIANDER, WAYMIRE 00, KAHANE, PEYRIÈRE 75, BARRAL, MANDELBROT 02.

LINEARISATION EFFECT: PICTURE

- TWO POWER-LAWS, TWO FUNCTIONS OF q :

- BARE CASCADE:

$$\mathbb{E}Q_r(t)^q = r^{\varphi(q)}, \quad q \in \mathcal{R}.$$

- DRESSED CASCADE:

$$\left. \begin{aligned} \mathbb{E}T_{Q_0}(t, a; \beta_0)^q &= c_q |a|^{\zeta(q)}, & q \in [q_c^-, q_c^+], \\ \mathbb{E}T_{Q_0}(t, a; \beta_0)^q &= \infty, & \text{ELSE,} \end{aligned} \right\}$$

WITH:

$$\left. \begin{aligned} \zeta(q) &= 1 + qh_*^-, & q \in [q_c^-, q_*^-], \\ \zeta(q) &= \varphi(q), & q \in [q_*^-, q_*^+], \\ \zeta(q) &= 1 + qh_*^+, & q \in [q_*^+, q_c^+]. \end{aligned} \right\}$$


- CONFUSION BETWEEN $\varphi(q)$ AND $\zeta(q)$:

- MULTIPLICATIVE CASCADE: $\varphi(q)$, $q \in \mathcal{R}$,

- SCALING EXPONENTS: $\zeta(q)$, $q \in [q_c^-, q_c^+]$.

LINEARISATION EFFECT: SKETCHED VIEWS

Moments




$E A_{\tau}(t)^q =$




Estimated $\zeta(q,n)$

EI



Estimated $\zeta(q,n)$

EII & EIII



LINEARISATION EFFECT: IMPACTS AND IMPORTANCE

CONSEQUENCES: RECAST THE USUAL GOALS :

- ESTIMATE THE INTEGRAL SCALE AND THE RESOLUTION OF THE CASCADE,
⇒ I.E., FIND A SCALING RANGE $[a_m, a_M]$
- ESTIMATE THE CRITICAL PARAMETERS $D_*^\pm, h_*^\pm, q_*^\pm$,
- ESTIMATE THE $\zeta(q)$ FOR $q \in [q_*^-, q_*^+]$,
→ VISIT B. LASHERMES'S POSTER.



IMPORTANCE OF THE LINEARISATION EFFECT:

- DISCRIMINATION OF MF MODELS BASED ON $\hat{\zeta}(q, n)$,
- DISCRIMINATION BETWEEN MONOFRACTAL AND MULTIFRACTAL,

NEGATIVE VALUES OF q

DIFFICULTIES ?


- FINITENESS ? $S_q(j) = (1/n_j) \sum_{k=1}^{n_j} |d_X(j, k)|^q < \infty$?
- NUMERICAL INSTABILITY ? $d_X(j, k) \simeq 0 \rightarrow |d_X(j, k)|^q = \infty$
- THEORY ? FULL MULTIFRACTAL SPECTRUM



SOLUTIONS ?

NEGATIVE VALUES OF q S - SOLUTION 1

AGGREGATION: $T_X(a, t) = \frac{1}{aT_0} \int_t^{t+aT_0} X(u) du$




APPLIES ONLY TO POSITIVE DATA (MEASURE)

NEGATIVE VALUES OF q S - SOLUTION 2



WT MODULUS MAXIMA (ARNEODO ET AL.)

$$L_X(a, t_k) = \sup_{a' < a} |T_X(a', t_k(a'))|$$

WTMM



Skeleton of the Wavelet Transform





COMPUTATIONALLY EXPENSIVE

NEGATIVE VALUES OF q S - SOLUTION 3

WAVELET LEADERS: (JAFFARD ET AL.)

$$d_X(j, k) \rightarrow L_X(j, k) = \sup_{j' < j} d_X(j', 2^{-j'})$$



COMPUTATIONALLY EFFICIENT AND EXCELLENT STATISTICAL PERFORMANCE

BEYOND POWER LAWS

- SELF-SIMILARITY:

$$\mathbb{E}|d_X(j, k)|^q = C_q(2^j)^{qH} = C_q \exp(qH \ln 2^j)$$

- POWER LAWS,
- $\forall 2^j$ (FOR ALL SCALES),
- $\forall q / \mathbb{E}|d_X(j, k)|^q < \infty$,
- A SINGLE PARAMETER H
- ADDITIVE STRUCTURE.

- MULTIFRACTAL

$$\mathbb{E}|d_X(j, k)|^q = C_q(2^j)^{\zeta(q)} = C_q \exp(\zeta(q) \ln 2^j)$$

- POWER LAWS,
- $\forall 2^j < L$, (FOR FINE SCALES ONLY, IN THE LIMIT $2^j \rightarrow 0$),
- $\forall q$?
- A WHOLE COLLECTION OF SCALING PARAMETER $\zeta(q)$
- MULTIPLICATIVE STRUCTURE.

- BEYOND POWER LAWS : WARPED INF. DIV. CASCADES

$$\mathbb{E}|d_X(j, k)|^q = C_q(2^j)^{qH} = C_q \exp(qH \ln 2^j)$$

$$\mathbb{E}|d_X(j, k)|^q = C_q(2^j)^{\zeta(q)} = C_q \exp(\zeta(q) \ln 2^j)$$

$$\mathbb{E}|d_X(j, k)|^q = \quad \quad \quad = C_q \exp(\zeta(q)n(2^j))$$

→ VISIT PIERRE CHAINAIS'S POSTER

CONCLUSIONS AND REFERENCES

ANALYSING SCALING IN DATA ?

- THINK WAVELET
 - EFFICIENCY,
 - PRACTICAL AND CONCEPTUAL ADEQUATION AND SIMPLICITY,
 - ROBUSTNESS AGAINST NON STATIONARITIES,
 - EASY TO USE, LOW COAST, REAL TIME ON LINE.

MODELLING SCALING IN DATA ?

- THINK SELF SIMILARITY VERSUS MULTIPLICATIVE CASCADES,
- AND POSSIBLY ADD LONG MEMORY.
- ALSO SCALING MAY NOT BE POWER LAWS

REFERENCES AND RESOURCES, VISIT :

- perso.ens-lyon.fr/patrice.abry
- inrialpes.fr/is2/~pgoncalv
- www.cubinlab.ee.mu.oz.au/~darryl
- fraclab
- www.isima.fr/~chainais